Fresh start. Cool possibilities.

Is reducing stubborn fat one of your New Year's goals? Schedule a consultation to see if freezing away stubborn fat with the CoolSculpting® treatment is right for you.





FREEZE AWAY STUBBORN FAT WITH COOLSCULPTING® AND GET STARTED ON YOUR RESOLUTION GOALS


- Non-surgical
- Little or no downtime
- FDA-cleared, safe and proven-effective

Call today to schedule your FREE consultation!



SUMAYAH J TALIAFERRO MD 404-873-1795

232 19th Street, Suite 7230
Atlanta, GA 30363
www.atlantadermatologyaesthetics.com





*CoolSculpting® is the treatment doctors use most for non-invasive fat removal.

Results and patient experience may vary.

Uses

The CoolSculpting® procedure is FDA-cleared for the treatment of visible fat bulges in the submental (under the chin) and submandibular (under the jawline) areas, thigh, abdomen and flank, along with bra fat, back fat, underneath the buttocks (also known as banana roll) and upper arm. It is also FDA-cleared to affect the appearance of lax tissue with submental area treatments. The CoolSculpting® procedure is not a treatment for weight loss.

Important Safety Information

The CoolSculpting® procedure is not for everyone. You should not have the CoolSculpting® procedure if you suffer from cryoglobulinemia, cold agglutinin disease, or paroxysmal cold hemoglobinuria.

Tell your doctor if you have any medical conditions including recent surgery, pre-existing hernia, and any known sensitivities or allergies.

During the procedure you may experience sensations of pulling, tugging, mild pinching, intense cold, tingling, stinging, aching, and cramping at the treatment site. These sensations subside as the area becomes numb. Following the procedure, typical side effects include temporary redness, swelling, blanching, bruising, firmness, tingling, stinging, tenderness, cramping, aching, itching, or skin sensitivity, and sensation of fullness in the back of the throat after submental or submandibular area treatment.

Rare side effects may also occur. CoolSculpting® may cause a visible enlargement in the treated area which may develop two to five months after treatment and requires surgical intervention for correction.

Please see full Important Safety Information at coolsculpting.com

